



# Earth Science with the Stratospheric Aerosol and Gas Experiment III (SAGE III) on the International Space Station



Joe Zawodny, Jean-Paul Vernier,  
Larry Thomason, Marilee Roell ,  
Mike Pitts, Randy Moore,  
Charles Hill, **David Flittner**, Rob  
Damadeo, Mike Cisewski



# SAGE III Science Objectives

NEED – enhance our understanding of ozone recovery and climate change processes in the upper atmosphere

HOW – monitor the vertical distribution of aerosol, ozone, and other trace gases in the Earth's stratosphere and troposphere


## **SAGE III/ISS provides data to:**

- Assess the recovery in the distribution of ozone
- Extend aerosol measurement records needed for climate and ozone models
- Gain further insight into key processes contributing to ozone and aerosol variability



# Ozone is Central to the Stratosphere


- Stratospheric ozone screens-out biologically harmful UV-C & UV-B sunlight.
- Ozone absorption of sunlight produces thermal structure of the atmosphere.





# Stratospheric Science Needs

## Stratospheric Chemistry & Climate Interactions




- This simplified schematic illustrates the parameters and process that control ozone
- Ozone Depleting Substances and Green House Gases can be measured from the ground as long as the dynamics can be modeled
- Ozone and Aerosol Profiles need to be measured
- Trends in Temperature and Water Vapor are inadequately measured



# Stratospheric Science Results

- The multi-decadal SAGE data are the international standard for ozone and aerosol.
- SAGE III predecessors have documented the effectiveness of the Montreal Protocol ban on Ozone Depleting Substances.
- Stratospheric aerosol time series is a vital component to understanding ozone changes.




*Yang et al., 2006*





# Model Ozone Uncertainties


35-60N Annual Mean Total Column Ozone Relative to 1980



- Ozone loss varies greatly with altitude and latitude
- Model estimates of the loss and eventual recovery differ
- Aggregate uncertainty due to modeling processes, and future ODS & GHG changes
- 2015-2020 measurements of ozone will improve understanding




# Stratospheric Aerosol Layer






# Stratospheric Aerosol Forcing Must be Included to Understand and Predict Climate Change

- Isolated colossal volcanic eruptions have significant cooling for a limited time.




- Increased background loading during 2000-2010 likely cause of global warming slow-down (Solomon, 2011).








# Measurement Strategy




Solar Occultation Geometry






# SAGE III Instrument Features

- A UV-Vis-NIR spectrometer
- Multiple modes of operation
  - Solar, lunar, limb scatter
- Surface/cloud top to 50 km, <1 km vertical resolution
- 87 channels (~1-3-nm resolution) between 280 and 1040 nm in solar occultation mode
- 64 kg, 102 watts, 0.12 Mbps






# Spectral Sampling





# Instrument Payload: NASA SMD, HEOMD & ESA contributions



NOTE: New hardware in **blue**




# ISS Orbit is Ideal!



Solar  
Lunar

Occultation Coverage

Alt	Inc	RA
350	51.6°	0°
350	51.6°	0°




	Rise	Set
Sun	◇	□
Moon	+	x






# SAGE III Climate Continuity Mission is empowered by NASA SMD & HEOMD, and ESA





# Launch Configuration


Manifest: February, 2016







# SAGE III on ISS, an Earth Science Mission on the International Space Station



**ELC-4**



# Summary

---

- The SAGE series has a long heritage and history of delivering outstanding and unique science products.
- SAGE III/ISS is a climate continuity mission addressing critical science needs.
- The ISS is an exceptional national asset in an ideal orbit for SAGE III to contribute internationally.
- SAGE III/ISS is designed to meet the core science objectives, while capturing data for additional science discoveries.